

FINITE REPRESENTATIONS OF A QUIVER ARISING FROM STRING THEORY

XINYUN ZHU

ABSTRACT. Inspired by Cachazo, Katz and Vafa (“Geometric transitions and $\mathcal{N} = 1$ quiver theories” (hep-th/0108120)), we examine representations of “ $N = 1$ quivers” arising from string theory. We derive some mathematical consequences of the physics, and show that these results are a natural extension of Gabriel’s ADE theorem. Extending the usual ADE case that relates quiver representations to curves on surfaces, we relate these new quiver representations to curves on threefolds.

1. INTRODUCTION

In 1972, Gabriel [13] published his celebrated theorem about the finite representation type of ADE quivers without relations. Since then the study of finite representation type quiver theory has been an important topic because it provides a successful way to solve problems in the representation theory of algebras and Lie groups. Recently, it has attracted the attention of physicists, (see [3]) due to its close relation with the study of D-branes. A special type of quiver arising from string theory, which we will call an “ $N = 1$ ADE quiver”, was introduced in [3](see Definition 3.1.1). This $N = 1$ ADE quiver has a close relation with the usual ADE quiver. The representations of $N = 1$ ADE quivers will satisfy the following important relations (3.2),

$$\sum_i s_{ij} Q_{ji} Q_{ij} + P'_j(\Phi_j) = 0, \quad Q_{ij} \Phi_j = \Phi_i Q_{ij} = 0.$$

The purpose of this paper is to construct, under certain conditions, a finite-to-one correspondence between the simple representations of an $N = 1$ ADE quiver and the positive roots of the usual ADE quiver; this matches the physicists’ predictions.

The “reflection functors” which were used in [11] to reprove Gabriel’s Theorem provide us a way to attack this problem. In this paper, we first modify the “reflection functors” in [11], and then apply our modified

Date: February 2, 2008.

reflectors to get our Main Theorem in Section 4.2. Some related results using different methods were given in [17].

This paper is organized as follows. In Section 2, we briefly review the the geometry of threefolds with small resolutions. In Section 3, we give the definition of $N = 1$ ADE quivers and their representations, and we introduce our modified reflection functors. In Section 4, we apply our modified reflection functors to get a proof of the Main Theorem in Section 4.2. In Section 5, I give a correspondence between indecomposable representations and ADE configuration of curves.

Acknowledgements. The contents of this article derives from a part of my doctoral dissertation. I wish to thank my thesis adviser, Professor Sheldon Katz, for his help and encouragement. It was an honor and a privilege to have been his student, and I will be forever grateful for his mathematical assistance and inspiration, moral and financial support, unfailing kindness and inexhaustible patience.

2. A QUICK REVIEW OF THE GEOMETRY OF THREEFOLDS FOR A GENERAL ADE FIBRATION OVER $\mathbf{C}$.

In this section, we refer the reader to [3] and [10]. Let $C \subset Y$ be a rational curve (not necessarily irreducible) in a 3-fold Y with K_Y trivial in a neighborhood of C and $\pi : Y \rightarrow X$ a birational morphism such that $\pi(C) = p \in X$ and $\pi|_{Y-C}$ is an isomorphism onto $X - p$. We consider an analytic neighborhood of p (still denoted X) and its inverse image under π (still denoted Y). By a lemma of Reid [16] (1.1, 1.14), the generic hyperplane section through p is a surface X_0 with an isolated rational double point, and the proper transform of X_0 is a partial resolution $Y_0 \rightarrow X_0$ (i.e. the minimal resolution $Z_0 \rightarrow X_0$ factors through $Y_0 \rightarrow X_0$).

The partial resolution $Y_0 \rightarrow X_0$ determines combinatorial data $\Gamma_0 \subset \Gamma$ consisting of an ADE Dynkin diagram Γ (the type of the singularity p) and a subgraph Γ_0 (the dual graph of the exceptional set of Y_0).

Let $\mathcal{Z} \rightarrow \text{Def}(Z_0)$, $\mathcal{Y} \rightarrow \text{Def}(Y_0)$, and $\mathcal{X} \rightarrow \text{Def}(X_0)$ be semi-universal deformations of Z_0 , Y_0 , and X_0 . Following [10], there are identifications

$$(2.1) \quad \text{Def}(Z_0) \cong V =: \text{Res}(\Gamma)$$

$$(2.2) \quad \text{Def}(Y_0) \cong V/\mathfrak{W}_0 =: \text{PRes}(\Gamma, \Gamma_0)$$

$$(2.3) \quad \text{Def}(X_0) \cong V/\mathfrak{W} =: \text{Def}(\Gamma)$$

where V is the complex root space associated to Γ and $\mathfrak{W}$ is its Weyl group. $\mathfrak{W}_0 \subset \mathfrak{W}$ is the subgroup generated by reflections of the simple roots corresponding to $\Gamma - \Gamma_0$. Deformations of Z_0 or Y_0 can be

blown down to give deformations of X_0 ([19] Theorem 1.4) and the induced classifying maps are given by the natural maps $V \rightarrow V/\mathfrak{W}$ and $V/\mathfrak{W}_0 \rightarrow V/\mathfrak{W}$ under the above identifications.

We can view X as the total space of a 1-parameter family X_t defined by the classifying map

$$g : \Delta \rightarrow \text{Def}(\Gamma).$$

where $\Delta \subset \mathbb{C}$ is a small disk. Similarly, we get the compatible family Y_t given by a map

$$f : \Delta \rightarrow \text{PRes}(\Gamma, \Gamma_0).$$

That is, we get the diagram

$$\begin{array}{ccccc} \mathcal{Z} & \xrightarrow{\tilde{\sigma}} & \mathcal{Y} & \xrightarrow{\tilde{\rho}} & \mathcal{X} \\ \downarrow & & \downarrow & & \downarrow \\ \text{Res}(\Gamma) & \xrightarrow{\sigma} & \text{PRes}(\Gamma, \Gamma_0) & \xrightarrow{\rho} & \text{Def}(\Gamma) \\ & & \uparrow f & \nearrow g & \\ & & \Delta & & \end{array}$$

By [10], $\mathcal{Y}$ is a blowup of $\mathcal{X} \times_{V/\mathfrak{W}} V/\mathfrak{W}_0$ and $\mathcal{Z}$ is a blowup of $\mathcal{X} \times_{V/\mathfrak{W}} V$. By construction, Y is the pullback of $\mathcal{Y}$ by f and X is the pullback of $\mathcal{X}$ by g .

2.1. The geometry of threefolds with small resolutions for a general ADE fibration over $\mathbb{C}$. Let $X \subset \mathbb{C} \times \mathbb{C}^3$ be an ADE fibration over $\mathbb{C}$. Let $t_i : \mathbb{C} \rightarrow V$ be a map ($1 \leq i \leq n+1$ in A_n case, $1 \leq i \leq n$ in D_n and E_n case), where V is the complex root space defined on (2.1). We consider the A_n case first. Then $X \subset \mathbb{C} \times \mathbb{C}^3$ is defined by the equation

$$xy = z^{n+1} + \alpha_2(\omega)z^{n-1} + \cdots + \alpha_{n+1}(\omega).$$

We write $h : \mathbb{C} \rightarrow V \subset \mathbb{C}^{n+1}$ as

$$h = (t_1, \dots, t_{n+1}) : \mathbb{C} \rightarrow \mathbb{C}^{n+1}, \sum_{i=1}^{n+1} t_i = 0.$$

Referring to [10], $\alpha_1, \dots, \alpha_{n+1}$ are elementary symmetric functions in $t_i, \dots, t_{n+1}$.

Let Z be the closure of the graph of the rational map

$$X \rightarrow (\mathbb{P}^1)^n, (x, y, z, \omega) \rightarrow \left[x, \prod_{j=1}^i (z + t_j(\omega)) \right]_i.$$

and let (u_i, v_i) be coordinates of the i -th $\mathbb{P}^1$ in $(\mathbb{P}^1)^n$. Using the identities

$$[x, z + t_1(\omega)] = [(z + t_2(\omega)) \cdots (z + t_n(\omega)), -y],$$

we get

$$xv_j = u_j \prod_{i=1}^j (z + t_i(\omega)) \quad (1 \leq j \leq n),$$

and

$$\prod_{i=k+1}^j (z + t_i(\omega)) u_j v_k = u_k v_j \quad (1 \leq k < j \leq n).$$

We refer the reader to [10] for the more complicated fibrations of the D and E cases. We list the defining equation of their deformations as follows:

$$D_n : x^2 + y^2 z + \frac{\prod_{i=1}^n (z + t_i^2(\omega)) - \prod_{i=1}^n t_i^2(\omega)}{z} + 2 \prod_{i=1}^n t_i(\omega) y$$

$$E_6 : x^2 + z^4 + y^3 + \epsilon_2 y z^2 + \epsilon_5 y z + \epsilon_6 z^2 + \epsilon_8 y + \epsilon_9 z + \epsilon_{12}$$

$$E_7 : x^2 + y^3 + y z^3 + \epsilon_2 y^2 z + \epsilon_6 y^2 + \epsilon_8 y z + \epsilon_{10} z^2 + \epsilon_{12} y + \epsilon_{14} z + \epsilon_{18}$$

$$E_8 : x^2 + y^3 + z^5 + \epsilon_2 y z^3 + \epsilon_8 y z^2 + \epsilon_{12} z^3 + \epsilon_{14} y z + \epsilon_{18} z^2 + \epsilon_{20} y + \epsilon_{24} z + \epsilon_{30}$$

where ϵ_i are complicated homogeneous polynomials in the t'_j s of degree i and invariant under the permutation of the t'_j s. We define entire functions $\{P'_i(t)\}$ as follows,

$$(2.4) \quad A_n : P'_i = t_i - t_{i+1} \quad i = 1, \dots, n$$

$$(2.5) \quad D_n : P'_i = t_i - t_{i+1} \quad i = 1, \dots, n-1 \quad \text{and} \quad P'_n = t_{n-1} + t_n$$

$$(2.6) \quad E_n : P'_i = t_i - t_{i+1} \quad i = 1, \dots, n-1 \quad \text{and} \quad P'_n = -t_1 - t_2 - t_3$$

3. A DESCRIPTION OF N=1 ADE QUIVERS AND REFLECTION FUNCTORS

3.1. Describing N=1 ADE quivers. This requires some detailed explanation, mainly of the relations (3.1), which distinguish these from ADE quivers. To make our presentation intelligible to non-experts, we briefly recall some definitions and established facts. (Here all vectors are over a field k .)

A *quiver* $\Gamma = (V_\Gamma, E_\Gamma)$ —without relations—is a directed graph.

A *representation* (V, f) of a quiver Γ is an assignment to each vertex $i \in V_\Gamma$ of a vector space $V(i)$, and to each directed edge $ij \in E_\Gamma$ of a linear transformation $f_{ji} : V(i) \rightarrow V(j)$.

$$\begin{array}{ccc} V(i) & \xrightarrow{h_i} & V'(i) \\ f_{ji} \downarrow & & f'_{ji} \downarrow \\ V(j) & \xrightarrow{h_j} & V'(j) \end{array}$$


Definition 3.1.1. Given an *ADE* Dynkin diagram Γ ,

$$A_n : \quad 1 - 2 - 3 - \cdots - n$$


$$D_n : \quad \begin{array}{ccccccc} & & & & n & & \\ & & & & | & & \\ 1 & - & 2 & - & \cdots & - & (n-2) - (n-1) \end{array}$$

$$E_n : \quad \begin{array}{ccccccc} & & & n & & & \\ & & & | & & & \\ 1 & - & 2 & - & 3 & - & \cdots - (n-1) \end{array}$$

$$A_n : \quad \begin{array}{c} e_1 \quad e_2 \quad e_3 \quad \quad e_n \\ \curvearrowright \quad \curvearrowright \quad \curvearrowright \quad \quad \curvearrowright \\ 1 \xrightleftharpoons[e_{12}]{e_{21}} 2 \xrightleftharpoons[e_{23}]{e_{32}} 3 \quad \dots \quad n \end{array}$$



The associated directed graphs for E_n ($n = 6, 7, 8$) are,



Then the $N = 1$ ADE quivers are just the above associated directed graphs (3.1)

$$(3.1) \quad \sum_i s_{ij} e_{ji} e_{ij} + P'_j(e_j) = 0, \quad e_{ij} e_j = e_i e_{ij}$$

where

$$\begin{cases} s_{ij} = 0 & \text{if } i \text{ and } j \text{ are not adjacent} \\ s_{ij} = 1 & \text{if } i \text{ and } j \text{ are adjacent and } i > j \\ s_{ij} = -1 & \text{if } i \text{ and } j \text{ are adjacent and } i < j \end{cases}$$

and where $P'_j(x)$ is a certain fixed polynomial $\forall j$.

If (V, f) is a representation of an $N = 1$ ADE quiver, the corresponding structures are

$$A_n : \quad \begin{array}{c} \Phi_1 \\ \curvearrowright \\ V(1) \end{array} \xrightleftharpoons[Q_{12}]{Q_{21}} \begin{array}{c} \Phi_2 \\ \curvearrowright \\ V(2) \end{array} \xrightleftharpoons[Q_{23}]{Q_{32}} \begin{array}{c} \Phi_3 \\ \curvearrowright \\ V(3) \end{array} \quad \cdots \quad \begin{array}{c} \Phi_n \\ \curvearrowright \\ V(n) \end{array}$$

$$\begin{array}{c}
D_n : \quad \begin{array}{c} \begin{array}{ccc} \overset{\Phi_1}{\curvearrowright} & & \overset{\Phi_2}{\curvearrowright} \\ V(1) & \xrightleftharpoons[Q_{12}]{Q_{21}} & V(2) \end{array} & \cdots & \begin{array}{ccc} \overset{\Phi_{n-2}}{\curvearrowright} & & \overset{\Phi_{n-1}}{\curvearrowright} \\ V(n-2) & \xrightleftharpoons[Q_{n-2,n-1}]{Q_{n-1,n-2}} & V(n-1) \end{array} \\ & & \begin{array}{c} \begin{array}{c} \uparrow \downarrow \\ Q_{n-2,n} \end{array} \\ V(n) \\ \begin{array}{c} \downarrow \uparrow \\ Q_{n,n-2} \end{array} \end{array} \end{array} \\
\\
E_n : \quad \begin{array}{c} \begin{array}{cccc} \overset{\Phi_1}{\curvearrowright} & \overset{\Phi_2}{\curvearrowright} & \overset{\Phi_3}{\curvearrowright} & \overset{\Phi_4}{\curvearrowright} \\ V(1) & \xrightleftharpoons[Q_{12}]{Q_{21}} & V(2) & \xrightleftharpoons[Q_{23}]{Q_{32}} & V(3) & \xrightleftharpoons[Q_{34}]{Q_{43}} & V(4) \end{array} & \cdots & \begin{array}{ccc} \overset{\Phi_{n-2}}{\curvearrowright} & & \overset{\Phi_{n-1}}{\curvearrowright} \\ V(n-2) & \xrightleftharpoons[Q_{n-2,n-1}]{Q_{n-1,n-2}} & V(n-1) \end{array} \\ & & \begin{array}{c} \begin{array}{c} \uparrow \downarrow \\ Q_{3n} \end{array} \\ V(n) \\ \begin{array}{c} \downarrow \uparrow \\ Q_{n3} \end{array} \end{array} \end{array}
\end{array}$$

where we have write $Q_{ij} = f_{e_{ij}}$, $\Phi_j = f_{e_j}$. And the relation (3.1) becomes

$$(3.2) \quad \sum_i s_{ij} Q_{ji} Q_{ij} + P'_j(\Phi_j) = 0, \quad Q_{ij} \Phi_j = \Phi_i Q_{ij}.$$

3.2. Reflection functors. Given an $N = 1$ ADE quiver Γ and $k \in V_\Gamma$, denote by Γ_k^+ the quiver defined by dropping all arrows starting from k , and denote by Γ_k^- the quiver defined from Γ by dropping all arrows ending at k .

Given a representation V of an $N = 1$ ADE quiver Γ , we can define a representation of Γ_k^+ which we still denote as V , by forgetting all maps which have domain $V(k)$. Similarly, we define a representation of Γ_k^- which we still denote by V , by forgetting all maps which has range $V(k)$. Then we can apply the reflection functor F_k^+ in [11] to the representation V of Γ_k^+ and apply the reflection functor F_k^- in [11] to the representation V of Γ_k^- .

In the following definition 3.2.1, we modify the reflection functors in [11] for the purpose of this paper.

Definition 3.2.1. Let Γ be an $N = 1$ ADE quiver and k a vertex of Γ . Let

$$\Gamma^k = \{i \mid i \text{ adjacent to } k\}$$

For a quiver representation W of Γ_k^+ , define a representation $F_k^+(W)$ of Γ_k^- by

$$(3.3) \quad F_k^+(W)(i) = \begin{cases} W(i) & \text{if } i \neq k \\ \ker h & \text{if } i = k \end{cases}$$

where

$$h : \oplus_{i \in \Gamma^k} W(i) \rightarrow W(k)$$

is defined by

$$h((x_i)_{i \in \Gamma^k}) = \sum_{i \in \Gamma^k} s_{ik} Q_{ki} x_i$$

If $i, j \neq k$, we define $Q'_{ij} = Q_{ij} : W(j) \rightarrow W(i)$. If $i \in \Gamma^k$, define $Q'_{ik} : F_k^+(W)(k) \rightarrow W(i)$ by

$$(3.4) \quad Q'_{ik}(x_j)_{j \in \Gamma^k} = -s_{ki} x_i$$

For a quiver representation U of Γ_k^- , define a representation $F_k^-(U)$ of Γ_k^+ by

$$(3.5) \quad F_k^-(U)(i) = \begin{cases} U(i) & \text{if } i \neq k \\ \text{coker } g & \text{if } i = k \end{cases}$$

where

$$g : U(k) \rightarrow \oplus_{i \in \Gamma^k} U(i)$$

is defined by

$$g(x) = (Q_{ik}x)_{i \in \Gamma^k}$$

and define $Q'_{ki} : U(i) \rightarrow F_k^-(U)(k)$ by the natural composition of

$$(3.6) \quad U(i) \rightarrow \oplus_{j \in \Gamma^k} U(j) \rightarrow F_k^-(U)(k)$$

Remark 3.2.1. The definitions of the $F_k^+(W)$ and Q'_{ik} in 3.2.1 are different than the corresponding definitions in [11], while $F_k^-(U)$ and Q'_{ki} in 3.2.1 are the same as the corresponding definitions in [11].

3.3. The action of the Weyl group on $\{P'_i\}$, $1 \leq i \leq n$. By [10], pp 461 and 463, we know the Weyl group $\mathfrak{W}_{A_n}$ of A_n is generated by reflections $r_1, \dots, r_n$, which act as permutations of $t_1, \dots, t_{n+1}$, where t_i is defined on Section 2.1.

In the A_n case, we can write $P'_i(x)$ in the relation given in (3.2) as

$$A_n : \quad P'_i = t_i - t_{i+1} \quad i = 1, \dots, n$$

Then the action of r_i extends to polynomials in the $\{t_i\}$. By [10], pp 461 and 463, we know the Weyl group $\mathfrak{W}_{D_n}$ of D_n is generated by reflections r_i , for $1 \leq i \leq n-1$, which act as permutations of $t_1, \dots, t_n$, together with r_n which is defined by

$$r_n(t_i) = \begin{cases} t_1 & \text{if } 1 \leq i \leq n-2 \\ -t_n & \text{if } i = n-1 \\ -t_{n-1} & \text{if } i = n \end{cases}$$

In the D_n case, we can write $P'_i(x)$ in the relation given in (3.2) as

$$D_n : \quad P'_i = t_i - t_{i+1} \quad i = 1, \dots, n-1$$

and

$$P'_n = t_{n-1} + t_n$$

Again, the action of r_i extends to polynomials in the $\{t_i\}$. By [10], pp 461 and 463, we know that the Weyl group $\mathfrak{W}_{E_n}$ of E_n is generated by reflections r_i for $1 \leq i \leq n-1$, which act as permutations of $t_1, \dots, t_n$, together with r_n , which is defined by

$$r_n(t_i) = \begin{cases} t_i - \frac{2}{3}(t_1 + t_2 + t_3) & \text{if } 1 \leq i \leq 3 \\ t_i + \frac{1}{3}(t_1 + t_2 + t_3) & \text{if } 4 \leq i \leq n \end{cases}$$

In the E_n case, we can write $P'_i(x)$ in the relation given in (3.2) as

$$E_n : \quad P'_i = t_i - t_{i+1} \quad i = 1, \dots, n \quad \text{and} \quad P'_n = -t_1 - t_2 - t_3$$

Once again, the action of r_i extends to polynomials in the $\{t_i\}$. Based on these definitions of r_i , $1 \leq i \leq n$, one can easily get the following Lemma 3.3.1.

Lemma 3.3.1. *Let $\mathfrak{W}_\Gamma$ be the Weyl group of the Dynkin diagram Γ and let $r_i \in \mathfrak{W}_\Gamma$ ($1 \leq i \leq n$) be a set of generators of reflections. If j is distinct from i and not adjacent to i , then $r_i(P'_j(\Phi_j)) = P'_j(\Phi_j)$. If j is adjacent to i and $j \neq i$, then $r_i(P'_j(\Phi_j)) = P'_j(\Phi_j) + P'_i(\Phi_j)$. Finally, $r_i(P'_i(\Phi_i)) = -P'_i(\Phi_i)$.*

4. FINITE-TO-ONE CORRESPONDENCE

In this section, we will give a proof, using reflection functors, that in the case of simple and distinct roots, the irreducible quiver representations are in finite-to-one correspondence with the contractible curves in the threefold.

4.1. Applying the reflection functors to $N = 1$ ADE quiver representations. Let Γ be an $N=1$ ADE quiver. Let

$$\mathcal{A}_\Gamma = \left\{ \sum_i n_i P'_i \mid n_i \in \mathbb{Z}, \text{ not all } n_i \text{ zero} \right\}$$

where P'_i , $1 \leq i \leq n$, are the polynomials in relation (3.2)

(*) Suppose no two elements $\sum n_i P'_i$, $\sum m_i P'_i$ of the set $\mathcal{A}_\Gamma$ have a common root unless there is a constant c with $m_i = cn_i$ for all i .

Lemma 4.1.1. *(*) holds for any generic collection of polynomials P'_i of positive degree.*

Proof. Left to the reader. $\square$

Lemma 4.1.2. *Let V be an $N = 1$ ADE quiver representation, let v_j be a λ -eigenvector of Φ_j . Then $Q_{ij}\Phi_j v_j$ is either a λ -eigenvector of Φ_i or 0.*

Proof. If v_j is an eigenvector of Φ_j corresponding to eigenvalue λ , then from (3.2), we get

$$Q_{ij}\Phi_j v_j = \Phi_i Q_{ij} v_j$$

which implies that

$$\lambda Q_{ij} v_j = \Phi_i Q_{ij} v_j$$

Hence, $Q_{ij} v_j$ is either an eigenvector of Φ_i corresponding to eigenvalue λ or a 0 vector. $\square$

Lemma 4.1.3. *Let V be a simple representation of an $N = 1$ ADE quiver. Then there exists λ such that if $v_i \in V(i) \neq 0$, then $\Phi_i v_i = \lambda v_i$.*

Proof. Let $\mathcal{A} = \{d | V(d) \neq 0\}$. Then $\mathcal{A}$ is connected. Otherwise, V is not simple. Let $a = \min \mathcal{A}$, then Φ_a has a eigenvector v_a with eigenvalue λ . For $l \in \mathcal{A}$, let $U(l)$ be the λ -eigenvector space of Φ_l . By Lemma 4.1.2, it's easy to see that $(W, g) = \{U(l) : l \in \mathcal{A}\}$ is a sub-representation of V . Since V is simple, $(W, g) = V$, which proves the result. $\square$

Therefore, to show that we have only finitely many simple representations, it suffices to consider representations V for which there exists a λ such that if $0 \neq v_d \in V(d)$, then $\Phi_d v_d = \lambda v_d$. In the rest of this section, we only consider quiver representations V with this property.

Lemma 4.1.4. *Let V be a simple representation of an $N = 1$ ADE quiver. Suppose V is not concentrated at vertex k . Then*

$$\dim (F_k^+(V))_k = \sum_{i \in \Gamma^k} \dim (V(i)) - \dim (V(k))$$

Proof. We know that $(F_k^+(V))(k) = \ker h$, where $h : \oplus_{i \in \Gamma^k} V(i) \rightarrow V(k)$ is defined by

$$h(x_i)_{i \in \Gamma^k} = \sum_{i \in \Gamma^k} s_{ik} Q_{ki} x_i$$

To prove the lemma is equivalent to prove that h is surjective.

Case I: $V(k) \neq 0$. If h is not surjective and $h \neq 0$, then we can replace $V(k)$ by $h(\oplus_{i \in \Gamma^k} V(i))$ to get a sub-representation of V . But this contradicts the simplicity of V . Case II: $V(k) = 0$. We get that h is surjective since $h \equiv 0$ in this case. $\square$

Lemma 4.1.5. *Let V be a simple representation of an $N = 1$ ADE quiver. Suppose V is not concentrated at vertex k . Then*

$$\dim (F_k^-(V))_k = \sum_{i \in \Gamma^k} \dim (V(i)) - \dim (V(k))$$

Proof. We know that $(F_k^-(V))(k) = \text{coker } g$, where $g : V(k) \rightarrow \bigoplus_{i \in \Gamma^k} V(i)$ is defined by $g(x) = (Q_{ik}x)_{i \in \Gamma^k}$. To prove the lemma is equivalent to prove that g is injective.

Case I: $V(k) \neq 0$. If $\ker g \neq 0$, then we can define a simple subrepresentation which concentrated at vertex k . This contradicts the simplicity of V .

Case II: $V(k) = 0$. We get that g is injective since $g \equiv 0$ in this case. $\square$

Lemma 4.1.6. *Let V be a simple representation of an $N = 1$ ADE quiver. Suppose V is not concentrated at vertex k . If $P'_k(\lambda) \neq 0$, then there is a natural isomorphism φ between $F_k^+(V)(k)$ and $F_k^-(V)(k)$.*

Proof. Let $g : V(k) \rightarrow \bigoplus_{i \in \Gamma^k} V(i)$ be defined by

$$g(x) = (Q_{ik}x)_{i \in \Gamma^k}$$

and $h : \bigoplus_{i \in \Gamma^k} V(i) \rightarrow V(k)$ be defined by

$$h(x_i)_{i \in \Gamma^k} = \sum_{i \in \Gamma^k} s_{ik} Q_{ki} x_i$$

We have

$$F_k^-(V)(k) = \text{coker } g$$

and

$$F_k^+(V)(k) = \ker h$$

Since V is simple and not concentrated at k , g is injective and h is surjective. We have

$$\dim F_k^+(V)(k) = \dim F_k^-(V)(k) = \sum_{i \in \Gamma^k} \dim V(i) - \dim V(k)$$

$$\begin{array}{ccccccc}
& & 0 & & & & \\
& & \downarrow & & & & \\
& & F_k^+(V)(k) & & & & \\
& & \downarrow h' & \searrow \varphi & & & \\
0 \longrightarrow & V(k) & \xrightarrow{g} & \oplus_{i \in \Gamma^k} V(i) & \xrightarrow{g'} & F_k^-(V)(k) & \longrightarrow 0 \\
& \searrow -P'_k(\lambda) & & \downarrow h & & & \\
& & & V(k) & & & \\
& & & \downarrow & & & \\
& & & 0 & & &
\end{array}$$

Since $P'_k(\lambda) \neq 0$, $\text{im } g \cap F_k^+(V)(k) = \{0\}$. Let $g' : \oplus_{i \in \Gamma^k} V(i) \rightarrow F_k^-(V)(k)$ be the natural surjective map induced by g and let $h' : F_k^+(V)(k) \rightarrow \oplus_{i \in \Gamma^k} V(i)$ be the natural inclusion map induced by h . Then $\varphi = g' \circ h' : F_k^+(V)(k) \rightarrow F_k^-(V)(k)$ is a natural isomorphism (Since $\dim F_k^+(V)(k) = \dim F_k^-(V)(k)$ and φ is injective by $\text{im } g \cap F_k^+(V)(k) = \{0\}$.) $\square$

Definition 4.1.1. By Lemma 4.1.6, if $P'_k(\lambda) \neq 0$, we can construct a new representation $F_k(V)$ of Γ by

$$F_k(V)(i) = \begin{cases} V(i) & \text{if } i \neq k \\ F_k^+(V)(k) & \text{if } i = k \end{cases}$$

defining Q'_{lk} as it is defined in $F_k^+(V)(k)$ and defining Q'_{km} as the composition map $P'_k(\lambda) \cdot \varphi^{-1} \circ Q'_{km} : V(m) \rightarrow F_k^+(V)(k)$, where $Q'_{km} : V(m) \rightarrow F_k^-(V)(k)$ is the natural map defined in $F_k^-(V)$ and $\varphi : F_k^+(V)(k) \rightarrow F_k^-(V)(k)$ is the isomorphism defined in Lemma 4.1.6.

If V is simple, we define

$$\Phi'_i : F_k(V)(i) \rightarrow F_k(V)(i)$$

by $\Phi'_i(x) = \lambda x$, where λ is the eigenvalue of Φ on $V(i)$ that appeared in the representation of V . Abusing notation, we still denote Φ'_i as Φ_i .

Lemma 4.1.7. *If V is a simple representation of $N = 1$ ADE quiver, then*

$$\sum_i \dim(V(i)) \cdot P'_i(\lambda) = 0$$

Proof. This follows from the fact that $\forall$ pair i and j , $\text{Tr } Q_{ij}Q_{ji} = \text{Tr } Q_{ji}Q_{ij}$, and $\forall, i$, $\text{Tr } \Phi_i = \lambda \cdot \dim V(i)$, where λ is an eigenvalue for

all Φ_i . Now take trace operation to relations (3.2) and then sum the resulting equations. The result follows. $\square$

Lemma 4.1.8. *Let V be a simple representation of an $N = 1$ ADE quiver Γ , not concentrated at vertex k . Then*

$$\sum \dim (F_k(V)) (i) r_k (P'_i(\lambda)) = \sum \dim V(i) P'_i(\lambda)$$

Proof.

$$\begin{aligned} & \sum \dim (F_k(V)) (i) r_k (P'_i(\lambda)) \\ = & \sum_{i \in \Gamma^k} \dim V(i) (P'_k(\lambda) + P'_i(\lambda)) + \sum_{j \in \Gamma - \Gamma^k} \dim V(j) P'_j(\lambda) \\ & + \left(-\dim V(k) + \sum_{i \in \Gamma^k} \dim V(i) \right) (-P'_k(\lambda)) \\ = & \sum \dim V(i) P'_i(\lambda) \end{aligned}$$

$\square$

Proposition 4.1.1. *Let V be a simple representation of an $N = 1$ ADE quiver which is not concentrated at vertex k . If $P'_k(\lambda) \neq 0$, then $F_k(V)$ satisfies the following new relations*

$$(4.1) \quad \sum_i s_{ij} Q'_{ji} Q'_{ij} + r_k (P'_j(\Phi_j)) = 0, \quad Q'_{ij} \Phi_j = \Phi_i Q'_{ij}.$$

Proof. If $i \notin \Gamma^k$ and $i \neq k$, where i is a vertex of Γ such that $V(i) \neq 0$, there is nothing to prove. For $j \in \Gamma^k \cup \{k\}$, $b \in (F_k(V))_j$, we have

$$Q'_{ij} \Phi_j b = \lambda Q'_{ij} b = \Phi_i Q'_{ij} b$$

For $i \in \Gamma^k$ and $x \in V(i)$, by Definition 4.1.1, we know that

$$Q'_{ki} x = P'_k(\lambda) \cdot \varphi^{-1} \circ \underline{Q'_{ki}} x$$

where $\underline{Q'_{ki}} x = [(x_j)_{j \in \Gamma^k}] \in F_k^-(V)(k)$, and

$$x_j = \begin{cases} 0 & \text{if } j \neq i \\ x & \text{if } j = i \end{cases}$$

After a short computation, we see that

$$Q'_{ki} x = (y_j)_{j \in \Gamma^k}$$

where

$$y_j = \begin{cases} P'_k(\lambda) x + s_{ik} Q_{ik} Q_{ki} x & \text{if } j = i \\ Q_{jk} s_{ik} Q_{ki} x & \text{if } j \neq i \end{cases}$$

It follows that

$$s_{ki}Q'_{ik}Q'_{ki}x = s_{ki}Q'_{ik}(y_j)_{j \in \Gamma^k} = -P'_k(\lambda)x - Q_{ik}s_{ik}Q_{ki}x$$

Hence for $i \in \Gamma^k$ we have

$$\begin{aligned} & \sum_j s_{ji}Q'_{ij}Q'_{ji}x + r_k(P'_i(\lambda))x \\ &= \sum_j s_{ji}Q'_{ij}Q'_{ji}x + r_k(P'_i(\lambda))x \\ &= \sum_j s_{ji}Q'_{ij}Q'_{ji}x + P'_i(\lambda)x + P'_k(\lambda)x \\ &= \sum_{j \neq k} s_{ji}Q_{ij}Q_{ji}x + s_{ki}Q'_{ik}Q'_{ki}x + P'_i(\lambda)x + P'_k(\lambda)x \\ &= \sum_{j \neq k} s_{ji}Q_{ij}Q_{ji}x - P'_k(\lambda)x - Q_{ik}s_{ik}Q_{ki}x + P'_i(\lambda)x + P'_k(\lambda)x \\ &= 0 \end{aligned}$$

Let $(x_i)_{i \in \Gamma^k} \in F_k^+(V)(k)$. Then

$$s_{ik}Q'_{ki}Q'_{ik}(x_i)_{i \in \Gamma^k} = Q'_{ki}x_i = (x_{i_j})_{j \in \Gamma^k}$$

where

$$x_{i_j} = \begin{cases} P'_k(\lambda)x_i + Q_{ik}s_{ik}Q_{ki}x_i & \text{if } j = i \\ Q_{jk}s_{ik}Q_{ki}x_i & \text{if } j \neq i \end{cases}$$

Hence we have

$$\begin{aligned} & \sum_{i \in \Gamma^k} s_{ik}Q'_{ki}Q'_{ik}(x_i)_{i \in \Gamma^k} + r_k(P'_k(\lambda))(x_i)_{i \in \Gamma^k} \\ &= \sum_{i \in \Gamma^k} s_{ik}Q'_{ki}Q'_{ik}(x_i)_{i \in \Gamma^k} - P'_k(\lambda)(x_i)_{i \in \Gamma^k} \\ &= \sum_{i \in \Gamma^k} (x_{i_j})_{j \in \Gamma^k} - P'_k(\lambda)(x_i)_{i \in \Gamma^k} \\ &= 0 \end{aligned}$$

□

Lemma 4.1.9. *If V is a simple representation of an $N = 1$ ADE quiver which is not concentrated at vertex k and if $(*)$ holds, then $F_k F_k(V) \cong V$. Consequently, $F_k(V)$ is a simple representation.*

Proof. We know that $Q'_{ki} : V(i) \rightarrow F_k(V)(k)$ is defined by $Q'_{ki}x_i = P'_k(\lambda)\varphi^{-1}\underline{Q}_{ki}x_i$, where $\underline{Q}_{ki} : V(i) \rightarrow F_k^-(V)(k)$ is the composition of $V(i) \rightarrow \oplus_{i \in \Gamma^k} V(i)$ and $\oplus_{i \in \Gamma^k} V(i) \rightarrow F_k^-(V)(k)$ (See Definition 4.1.1). We also know that

$$F_k F_k(V)(k) = \left\{ (x_i) \in \oplus_{i \in \Gamma^k} V(i) \mid \sum_{i \in \Gamma^k} s_{ik} Q'_{ki} x_i = 0 \right\}$$

We have

$$\sum_{i \in \Gamma^k} s_{ik} Q'_{ki} x_i = P'_k(\lambda) \varphi^{-1} \sum_{i \in \Gamma^k} s_{ik} \underline{Q}_{ki} x_i$$

Since $P'_k(\lambda) \neq 0$ and φ is an isomorphism, we get

$$F_k F_k(V)(k) = \{(-s_{ki} Q_{ik} x) \mid x \in V(k)\}$$

Let $g : V \rightarrow F_k F_k(V)$ be defined in the following way:

$$g_i = \begin{cases} i : V(i) \rightarrow F_k F_k(V)(i) = V(i) & \text{if } i \neq k \\ (-s_{ki} Q_{ik}) & \text{if } i = k \end{cases}$$

where $i : V(i) \rightarrow F_k F_k(V)(i) = V(i)$ is the identity map.

Then it is clear that (4.2) is commutative.

$$(4.2) \quad \begin{array}{ccc} V(k) & \xrightarrow{Q_{ik}} & V(i) \\ \downarrow g_k & & \downarrow g_i \\ F_k F_k(V)(k) & \xrightarrow{Q''_{ik}} & V(i) \end{array}$$

Let's check the commutativity of (4.3).

$$(4.3) \quad \begin{array}{ccc} V(i) & \xrightarrow{Q_{ki}} & V(k) \\ \downarrow g_i & & \downarrow g_k \\ V(i) & \xrightarrow{Q''_{ki}} & F_k F_k(V)(k) \end{array}$$

Let $(Q''_{ki}x_i)_j$ (resp. $(Q'_{ki}x_i)_j$) denote the j -th coordinate of $Q''_{ki}x_i$ (resp. $Q'_{ki}x_i$). We know that

$$(Q''_{ki}x_i)_j = \begin{cases} -P'_k(\lambda)x_i + Q'_{ik}s_{ik}Q'_{ki}x_i & \text{if } j = i \\ Q'_{jk}s_{ik}Q'_{ki}x_i & \text{if } j \neq i \end{cases}$$

where

$$(Q'_{ki}x_i)_j = \begin{cases} P'_k(\lambda)x_i + Q_{ik}s_{ik}Q_{ki}x_i & \text{if } j = i \\ Q_{jk}s_{ik}Q_{ki}x_i & \text{if } j \neq i \end{cases}$$

If $i > k$, then we have

$$\begin{aligned}
(Q''_{ki}x_i)_i &= -P'_k(\lambda)x_i + Q'_{ik}s_{ik}Q'_{ki}x_i \\
&= -P'_k(\lambda)x_i + P'_k(\lambda)x_i + Q_{ik}s_{ik}Q_{ki}x_i \\
&= Q_{ik}s_{ik}Q_{ki}x_i = Q_{ik}Q_{ki}x_i \\
&= Q_{ik}Q_{ki}x_i = -s_{ki}Q_{ik}Q_{ki}x_i
\end{aligned}$$

If $i > k$ and $j > k$, then we have

$$(Q''_{ki}x_i)_j = Q'_{jk}s_{ik}Q'_{ki}x_i = Q_{jk}s_{ik}Q_{ki}x_i = Q_{jk}Q_{ki}x_i = -s_{kj}Q_{jk}Q_{ki}x_i$$

If $i > k$ and $j < k$, then we have

$$(Q''_{ki}x_i)_j = Q'_{jk}s_{ik}Q'_{ki}x_i = Q'_{jk}Q'_{ki}x_i = -Q_{jk}Q_{ki}x_i = -s_{kj}Q_{jk}Q_{ki}x_i$$

If $i < k$, then we have

$$\begin{aligned}
(Q''_{ki}x_i)_i &= -P'_k(\lambda)x_i + Q'_{ik}h'_{ki}x_i \\
&= -P'_k(\lambda)x_i - Q'_{ik}Q'_{ki}x_i \\
&= -P'_k(\lambda)x_i + (P'_k(\lambda)x_i + Q_{ik}h_{ki}x_i) \\
&= Q_{ik}h_{ki}x_i = -Q_{ik}Q_{ki}x_i \\
&= -s_{ki}Q_{ik}Q_{ki}x_i
\end{aligned}$$

If $i < k$ and $j > k$, then we have

$$(Q''_{ki}x_i)_j = Q'_{jk}s_{ik}Q'_{ki}x_i = -Q'_{jk}Q'_{ki}x_i = -Q_{jk}s_{ik}Q_{ki}x_i = Q_{jk}Q_{ki}x_i = -s_{kj}Q_{jk}Q_{ki}x_i$$

If $i < k$ and $j < k$, then we have

$$(Q''_{ki}x_i)_j = Q'_{jk}s_{ik}Q'_{ki}x_i = -Q'_{jk}Q'_{ki}x_i = Q_{jk}s_{ik}Q_{ki}x_i = -Q_{jk}Q_{ki}x_i = -s_{kj}Q_{jk}Q_{ki}x_i$$

Therefore the diagram (4.3) is commutative.

Diagram (4.4) is commutative since λ is a common eigenvalue of $V(k)$ and $F_k F_k(V)(k)$.

$$\begin{array}{ccc}
(4.4) & V(k) & \xrightarrow{\lambda} V(k) \\
& \downarrow g_k & \downarrow g_k \\
& F_k F_k(V)(k) & \xrightarrow{\lambda} F_k F_k(V)(k)
\end{array}$$

We prove the later part of the Lemma here. Since V is not concentrated at vertex k , $\exists m \neq k$ such that $V(m) \neq 0$. It follows that $F_k(V)(m) = V(m) \neq 0$. Let $v \in F_k(V)(m)$ be a nonzero element. If $F_k(V)$ is not simple, then there exists, actually we can construct a simple sub-representation W of $F_k(V)$, such that $v \in W(m)$. It follows that

$F_k(W)$ is a proper sub-representation of $F_k F_k(V)$. Since $F_k F_k(V) \cong V$ and V is simple, this leads a contradiction. $\square$

Corollary 4.1.1. *Let Γ be an $N = 1$ ADE quiver. Let*

$$\mathcal{B}_\Gamma = \{r_i(P'_j(x)) | r_i \in \mathfrak{W}_\Gamma\}$$

where $\mathfrak{W}_\Gamma$ is the Weyl group of Γ and P'_j is the polynomial defined on relation (3.2). Suppose $(*)$ holds and each element in $\mathcal{B}_\Gamma$ has simple roots. If V is a simple representation, then either $F_k(V)$ is simple or $V \cong L_k$, where L_k is a simple representation concentrated at vertex k .

Proof. If V is simple and concentrated at vertex k , then $V \cong L_k$, where L_k is a simple representation concentrated at vertex k . Assume V is not concentrated at vertex k . Since V is simple, by Lemma 4.1.4 and Lemma 4.1.5, we can apply F_k to V . Then $F_k(V)$ is simple by the later part of Lemma 4.1.9. $\square$

4.2. A proof of the Main Theorem. Let Γ be a quiver. Following [11], for a representation V , we define $\dim(V) = (\dim V(i))_{i \in V_\Gamma}$. Denote by $\mathcal{C}_\Gamma = \{x = (x_\alpha) \mid x_\alpha \in \mathbb{Q}, \alpha \in V_\Gamma\}$, where $\mathbb{Q}$ denotes the set of rational numbers. We call a vector $x = (x_\alpha)$ positive (written $x > 0$) if $x \neq 0$ and $x_\alpha \geq 0$ for all $\alpha \in V_\Gamma$. For each $\beta \in V_\Gamma$, denote by σ_β the linear transformation in $\mathcal{C}_\Gamma$ defined by the formula $(\sigma_\beta x)_\gamma = x_\gamma$ for $\gamma \neq \beta$, $(\sigma_\beta x)_\beta = -x_\beta + \sum_{l \in \Gamma^\beta} x_l$, where $l \in \Gamma^\beta$ is the set of vertices adjacent to β .

For each vertex $\alpha \in V_\Gamma$ we denoted by Γ_α the set of edges containing α . Let Λ be an orientation of the graph Γ . We denote by $\sigma_\alpha \Lambda$ the orientation obtained from Λ by changing the directions of all edges $l \in \Gamma_\alpha$. Following [11], we say that a vertex i of a quiver Γ with orientation Λ is $(-)$ -accessible (resp. $(+)$ -accessible) if for any edge e having i as a vertex, we have the final vertex of $f(e)$ of e satisfying $f(e) \neq i$ (resp. the initial vertex $i(e)$ of e satisfying $i(e) \neq i$.) We say that a sequence of vertices $\alpha_1, \alpha_2, \dots, \alpha_k$ is $(+)$ -accessible with respect to Λ if α_1 is $(+)$ -accessible with respect to Λ , α_2 is $(+)$ -accessible with respect to $\sigma_{\alpha_1} \Lambda$, α_3 is $(+)$ -accessible with respect to $\sigma_{\alpha_2} \sigma_{\alpha_1} \Lambda$, and so on. We define a $(-)$ -accessible sequence similarly.

Definition 4.2.1. Let Γ be a graph without loops. We denote by $\mathcal{C}_\Gamma$ the linear space over $\mathbb{Q}$ consisting of sets $x = (x_\alpha)$ of rational numbers x_α ($\alpha \in \Gamma_V$). We call a vector $x = (x_\alpha)$ *positive* (written $x > 0$) if $x \neq 0$ and $x_\alpha \geq 0$ for all $\alpha \in \Gamma_V$. We denote by B the quadratic form on the space $\mathcal{C}_\Gamma$ defined by the formula $B(x) = \sum x_\alpha^2 - \sum_{l \in \mathcal{E}_\Gamma} x_{r_1(l)} x_{r_2(l)}$,

where $r_1(l)$ and $r_2(l)$ are the ends of the edge l . We denote by $<, >$ the corresponding symmetric bilinear form.

Lemma 4.2.1. [11, Lemma 2.3] *Suppose that the form B for the graph Γ is positive definite. Let $c = \sigma_n \cdots \sigma_2 \sigma_1$. If $x \in \mathcal{C}_\Gamma$, $x \neq 0$, then for some i the vector $c^i x$ is not positive.*

Main Theorem. *Let Γ be an $N = 1$ ADE quiver. Let $\mathcal{B}_\Gamma = \{r_i(P'_j(x))\}$, where $r_i \in \mathfrak{W}_\Gamma$ and P'_j , $j \in V_\Gamma$ are the polynomials defined in relation (3.2). Assume no element in $\mathcal{B}_\Gamma$ has a multiple root. If $(*)$ holds, then $N = 1$ ADE quivers have finite representation type.*

Proof. Let V be a simple representation of an $N = 1$ ADE quiver. Let $\mathcal{A} = \{i | V(i) \neq 0\}$. We can assume that $\mathcal{A}$ is connected. Otherwise, V would be decomposable. We apply the forgetful functors to V to get an $(+)$ -accessible (resp. $(-)$ -accessible) diagram (no loop)

$$\begin{array}{ccccccc}
 V(1) & \longleftarrow & \cdots & \longleftarrow & V(k) & \longleftarrow & \cdots & \longleftarrow & V(n) \\
 & & & & \uparrow & & & & \\
 & & & & V(l) & & & & \\
 & & & & \downarrow & & & & \\
 V(1) & \longrightarrow & \cdots & \longrightarrow & V(k) & \longrightarrow & \cdots & \longrightarrow & V(n) \\
 & & & & \downarrow & & & & \\
 & & & & V(l) & & & &
 \end{array}$$

(For the type A case, $V(l) = 0$.)

Let $c = \sigma_n \cdots \sigma_2 \sigma_1$. By [11], there exists k such that $c^k(\dim V) \not\geq 0$. By $(*)$, Lemma 4.1.7 and Lemma 4.1.8, we know that $\sum_i \dim V(i) \cdot P'_i(x)$ is the only element in $\mathcal{A}_\Gamma$ which vanishes at λ . By Corollary 4.1.1 and Proposition 4.1.1, this implies that there exist $\beta_1, \dots, \beta_l$ and a simple representation $L_{\beta_{l+1}}$ which is concentrated at a vertex of Γ such that

$$V = F_{\beta_1} \cdots F_{\beta_k}(L_{\beta_{k+1}})$$

V corresponds to the positive root

$$\dim V = \sigma_{\beta_1} \cdots \sigma_{\beta_k}(\overline{\beta_{k+1}})$$

where

$$\overline{(\beta_{k+1})} = \begin{cases} 0 & \text{if } i \neq k+1 \\ 1 & \text{if } i = k+1 \end{cases}$$

Since the usual ADE quiver only has finitely many positive roots, $N = 1$ ADE quivers have finite representation type. This finishes the proof of the theorem. $\square$

From the above Main Theorem, one can get the following Proposition 4.2.1.

Corollary 4.2.1. *Let Γ be an $N = 1$ ADE quiver. Let $\mathcal{B}_\Gamma = \{r_i(P'_j(x)) | r_i \in \mathfrak{W}_\Gamma\}$, where $\mathfrak{W}_\Gamma$ is the Weyl group of Γ and P'_j is the polynomial defined on relation (3.2). Assume each element in $\mathcal{B}_\Gamma$ has simple roots. If $(*)$ holds, then there is a finite-to-one correspondence between simple representations of $N = 1$ ADE quivers and the positive roots of ADE Dynkin diagram.*

Proof. We know that $\mathcal{B}_\Gamma$ has only finitely many elements. Each element of $\mathcal{B}_\Gamma$ which is in fact a polynomial has only finitely many simple roots. By our Main Theorem, each root of an element in $\mathcal{B}_\Gamma$ corresponds with a simple representation. Hence, the desired result follows. $\square$

5. A CORRESPONDENCE BETWEEN INDECOMPOSABLE REPRESENTATIONS AND ADE CONFIGURATION OF CURVES.

This section will give a correspondence between indecomposable representations of an $N=1$ ADE quiver and ADE configuration of curves.

To make the presentation easier to follow, we state the following Lemma from [16, (1.1),(1.14)] here.

Reid's Lemma. *Let $\pi: Y \rightarrow X$ be a resolution of an isolated Gorenstein threefold singularity $P \in X$. Suppose that the exceptional set of π has pure dimension 1. Let X_0 be a generic hyperplane section of X which passes through P . Then X_0 has a rational double point at P .*

Moreover, if X_0 is any hyperplane section through P with a rational double point, and Y_0 is its proper transform, then Y_0 is normal, and the minimal resolution $Z_0 \rightarrow X_0$ factors through the induced map $\pi|_{Y_0}: Y_0 \rightarrow X_0$.

The exceptional curve of the minimal resolution $Z_0 \rightarrow X_0$ is a rational curve whose dual graph is Γ . Thus the components of the exceptional divisor are rational curves C_{e_i} which are in one to one correspondence with the simple roots $e_1, \dots, e_n$ (n is the rank of the root system).

Let X be an ADE fibration with base $\mathbb{C}$, and Y be the small resolution of X . Let $\pi: Y \rightarrow X$ be the blow up map. An “ADE configuration of curves” in Y is a 1 dimensional connected projective scheme $C \subset Y$, such that

- (1) $\exists \bar{S} \subset Y, C \subset \bar{S}$
- (2) letting $S = \pi(\bar{S})$, then $\bar{S} \rightarrow S$ is a resolution of ADE singularities with exceptional scheme C .

We need the following proposition which is essentially part 3 of Theorem 1 in [10].

Proposition 5.0.1. *The irreducible components of the discriminant divisor $\mathfrak{D} \subset \text{Res}(\Gamma)$ are in one to one correspondence with the positive roots of Γ . Under the identification of $\text{Res}(\Gamma)$ with the complex root space U , the component $\mathfrak{D}_v$ corresponding to the positive root $v = \sum_{i=1}^n a_i e_i$ is $v^\perp \subset U$, ie the hyperplane perpendicular to v .*

Moreover, $\mathfrak{D}_v$ corresponds exactly to those deformations of Z_0 in $\mathcal{Z}$ to which the curve

$$C_v := \bigcup_{i=1}^n a_i C_{e_i}$$

lifts. For a generic point $t \in \mathfrak{D}_v$, the corresponding surface $\mathcal{Z}_t$ has a single smooth -2 curve in the class $\sum_{i=1}^n a_i [C_{e_i}]$ thus there is a small neighborhood B of t such that the restriction of $\mathcal{Z}$ to B is isomorphic to a product of $\mathbb{C}^{n-1}$ with the semi-universal family over $\text{Res}(A_1)$.

Let X be an ADE fibration with base $\mathbb{C}$, and Y be the small resolution of X .

The following example gives a concrete correspondence between the ADE configuration of curves in Y and the indecomposable representations of an N=1 ADE quiver in the A_2 case.

Example 5.0.1. Let X be defined by

$$A_2 : xy + (z + t_1(t))(z + t_2(t))(z + t_3(t)) = 0$$

with

$$t_1(t) + t_2(t) + t_3(t) = 0.$$

In the following Table 1, “Curve” means an ADE configuration of curves which is in the exceptional set of the fibration. I use “dim V ” to denote the dimension vector of an indecomposable representation of the N=1 ADE quiver.

Table 1 can be explained in the following way. If $t_1(\lambda) = t_2(\lambda) \neq t_3(\lambda)$ for some λ , then by Proposition 5.0.1, there exists an ADE configuration of curves $C \subset Y$. By [3], we know that $P'_1(\lambda) = t_1(\lambda) - t_2(\lambda) = 0$. Hence, by our Main Theorem, there exists an indecomposable representation V of an N=1 ADE quiver with $\dim V = (1, 0)$ that corresponds to C .

The other cases are similar, we omit them.

This example will be generalized in the following theorem.

Theorem 5.0.1. *Let X be a ADE fibration corresponding to Γ , with base $\mathbb{C}$. Let Y be a small resolution of X . Let $\mathcal{B}_\Gamma = \{r_i(P'_j(x)) | r_i \in \mathfrak{W}_\Gamma\}$,*


Condition	Singularity	Curve	$\dim V$
$t_1(\lambda) = t_2(\lambda) \neq t_3(\lambda)$	A_1	$\mathbb{P}^1$ 	$\begin{matrix} 1 & 0 \\ \bullet & \text{---} & \bullet \end{matrix}$
$t_1(\lambda) \neq t_2(\lambda) = t_3(\lambda)$	A_1	$\mathbb{P}^1$ 	$\begin{matrix} 0 & 1 \\ \bullet & \text{---} & \bullet \end{matrix}$
$t_1(\lambda) = t_3(\lambda) \neq t_2(\lambda)$	A_2		$\begin{matrix} 1 & 1 \\ \bullet & \text{---} & \bullet \end{matrix}$
$t_1(\lambda) \neq t_2(\lambda) \neq t_3(\lambda)$	-	-	-

TABLE 1.

where $\mathfrak{W}_\Gamma$ is the Weyl group of Γ and P'_j is the polynomial defined in relation (3.2). Assume no element in $\mathcal{B}_\Gamma$ has multiple roots and assume $(*)$ holds. Then there exists a 1-1 correspondence between the indecomposable representations of the $N = 1$ ADE quiver and the ADE configuration of curves in Y .

Proof. By Pinkham [14] and Katz-Morrison [10], we have the following commutative diagram

$$\begin{array}{ccc}
 Y & \longrightarrow & \mathcal{Y} \\
 \downarrow \pi & & \downarrow \\
 X & \longrightarrow & \mathcal{X} \\
 \downarrow \varphi & & \downarrow \\
 \mathbb{C} & \xrightarrow{f} & \mathbb{C}^n
 \end{array}$$

where $\mathbb{C}$ denotes the set of complex numbers and $\mathcal{Y}$ denotes the $\mathbb{C}^*$ -equivariant simultaneous resolution $\mathcal{Y} \rightarrow \mathcal{X}$ inducing $Y_0 \rightarrow X_0$. For an indecomposable representation V of the $N = 1$ ADE quiver Γ , we have

$$(5.1) \quad \sum \dim V(i) \cdot P'_i(\lambda) = 0$$

for some λ . The dimension vector $(\dim V(i))_{i \in V_\Gamma}$ will correspond to a positive root ρ . By (2.4), (2.5), and (2.6), we can express $P'_i(x)$, $i = 1, \dots, n$ in terms of t_i , $i = 1, \dots, n$. By Proposition 5.0.1 or part 3 of

Theorem 1 in Katz-Morrison [10, pp. 467], (5.1) will give an equation for $\rho^\perp$. Hence $f(\lambda) = (t_i(\lambda))_{i \in V_\Gamma} \in \rho^\perp$. It follows from Proposition 5.0.1 that there exists an ADE configuration of curves $C_\rho \subset \pi^{-1}(\lambda) \subset Y$.

Conversely, for an ADE configuration of curves $C \subset Y$, we have that $\varphi \circ \pi(C) = \lambda \in \mathbb{C}$ (Since π is projective, $\varphi \circ \pi(C)$ is projective in $\mathbb{C}$. It follows that $\varphi \circ \pi(C)$ is a finite subset of $\mathbb{C}$. Since C is connected, $\varphi \circ \pi(C)$ is connected in $\mathbb{C}$. Hence $\varphi \circ \pi$ is a point in $\mathbb{C}$.) Moreover, $\pi(C)$ is a point in X (By Katz-Morrison [10], we know that $\mathcal{X}$ is affine. Hence $\pi(C)$ is a point in X .) By Proposition 5.0.1, we know that $f(\lambda) \in \rho^\perp$ for some positive root ρ . Since we assume that each element in $\mathcal{B}_\Gamma$ has simple roots and $(*)$ holds, C corresponds to a unique positive root ρ . We can express ρ as $\rho = \sum a_i \cdot \rho_i$ where ρ_i is a simple positive root. From our Main Theorem, we can construct a simple representation V of $N = 1$ ADE quiver Γ which corresponds to the positive root ρ by applying the reflection functors. This finishes the proof of Theorem 5.0.1. □

REFERENCES

- [1] Maurice Auslander, Idun Reiten, and Sverre O. Smalø. *Representation theory of Artin algebras*. Cambridge studies in advanced mathematics ; 36. Cambridge University Press, 1997.
- [2] Bryan, Jim; Katz, Sheldon; Leung, Naichung Conan. Multiple covers and the integrality conjecture for rational curves in Calabi-Yau threefolds. *J. Algebraic Geom.*, 10(3):549–568., 2001.
- [3] F. Cachazo, S. Katz, and C. Vafa. Geometric transitions and $\mathcal{N}=1$ quiver theories, 2001. eprint hep-th/0108120.
- [4] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994.
- [5] Robin Hartshorne. *Algebraic geometry*. Graduate texts in mathematics, 52. New York : Springer-Verlag, 1977.
- [6] H.B.Laufer. On $\mathbb{CP}^1$ as exceptional set. In *Recent developments in several complex variables*, Ann. of Math. Stu. 100, pages 261–275. Princeton University Press, Princeton, 1981.
- [7] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [8] Jesús Jiménez. Contraction of nonsingular curves. *Duke Math. J.*, 65(2):313–332, 1992.
- [9] Sheldon Katz. Versal deformations and superpotentials for rational curves in smooth threefolds. In *Symposium in Honor of C. H. Clemens (Salt Lake City, UT, 2000)*, Contemp. Math., 312, pages 129–136. Amer. Math. Soc., Providence, RI, 2002.

- [10] Sheldon Katz and David R. Morrison. Gorenstein threefold singularities with small resolutions via invariant theory for Weyl groups. *J. Algebraic Geom.*, 1(3):449–530, 1992.
- [11] L.N.Bernstein, I.M.Gelfand, and V.A.Ponomarev. Coxeter functors and Gabriel’s theorem. *Usp.Mat.Nauk*, 28:17–32, 1973.
- [12] Hideyuki Matsumura. *Commutative ring theory*. London Mathematical Society Student Texts, 12. Cambridge University Press, Cambridge, 1988.
- [13] P.Gabriel. Unzerlegbare darstellungen I. *Manuscripta math*, 6:71–103, 1972.
- [14] Henry Pinkham. Factorization of birational maps in dimension 3. In *Singularities, Part 2 (Arcata, Calif., 1981)*, Proc. Sympos. Pure Math., 40, pages 343–371. Amer. Math. Soc., Providence, RI, 1983.
- [15] J. Le Potier. *Lectures on vector bundles*. Cambridge studies in advanced mathematics ; 54. Cambridge University Press, 1997.
- [16] Miles Reid. Minimal models of Canonical 3-folds. In *Algebraic varieties and analytic varieties*, Adv. Stud. Pure Math., 1, pages 131–180, Tokoyo, 1983. North-Holland, Amsterdam.
- [17] Balázs Szendrői. Sheaves on fibered threefolds and quiver sheaves. eprint math.AG/0506301.
- [18] Carlos T Simpson. Moduli of representations of the fundamental group of a smooth projective variety I. *Inst. Hautes Études Sci. Publ. Math.*, 79:47–129, 1994.
- [19] Jonathan M. Wahl. Equisingular deformations of the normal surface singularities. I. *Ann. of Math. (2)*, 104(2): 325–356, 1976.
- [20] Thomas Zerger. *Contracting rational curves on smooth complex threefolds*. PhD thesis, Oklahoma State University, 1996.
- [21] Thomas Zerger. Contraction criteria for reducible rational curves with components of length one in smooth complex threefolds. *Pacific J. Math*, 212(2):377–394, 2003.

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, STILLWATER, OK 74078

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA- CHAMPAIGN, URBANA, IL 61801

E-mail address: zhu5@math.uiuc.edu